

New multivariate tests for phylogenetic signal and trait correlations applied to ecophysiological phenotypes of nine *Manglietia* species

Li Zheng¹, Anthony R. Ives^{*2}, Theodore Garland Jr³, Bret R. Larget⁴, Yang Yu⁵ and Kunfang Cao^{*1}

¹Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming 650223, China; ²Department of Zoology, UW-Madison, Madison, Wisconsin 53706, USA; ³Department of Biology, University of California, Riverside, Riverside, California 92521, USA; ⁴Departments of Botany and Statistics, UW-Madison, Madison, Wisconsin 53706, USA; and ⁵Yunnan Provincial Environmental Protection Department, Appraisal Center for Environment and Engineering, 27 Xiyuannan Road, Kunming 650032, China

Summary

1. Phylogenetic signal – the similarity in trait values among phylogenetically related species – is pervasive for most types of traits in most organisms. Traits can often be categorized *a priori* into groups based on the level of biological organization, functional relations, developmental origins, or genetic underpinnings. Traits within such groups are often expected to be correlated and hence show similar levels of phylogenetic signal.

2. We developed multivariate statistical methods to test for phylogenetic signal in groups of traits while also incorporating estimates of trait measurement error (including within-species variation) that can obscure phylogenetic signal. Simultaneously, these methods produce estimates of correlations between traits that are corrected for phylogenetic relationships among species.

3. We applied these methods to data for 13 morphological and physiological traits gathered in a common-garden study of nine species of *Manglietia* (Magnoliaceae). The 13 traits fell into four groups: three traits involved photosynthesis [maximum net photosynthesis (A_{\max}), light saturation point (LSP), light compensation point]; three described leaf morphology (thickness of leaves, palisade tissue, sponge tissue); four related to plant growth (basal stem diameter, crown volume, leaf area, relative growth rate); and three measured thermal tolerance [critical temperature (T_{ch}), peak temperature (T_{\max}), temperature of half-inactivation (T_{50})]. We also constructed a molecular phylogeny for these species from 219 AFLP markers via maximum likelihood estimation under the assumption of sequential binary changes in DNA sequences.

4. Of the 13 traits, only two photosynthesis traits (A_{\max} and LSP) exhibited statistically detectable phylogenetic signal ($P < 0.05$) when analysed separately, whether using previously published univariate tests or our new univariate tests that incorporate measurement error. In contrast, multivariate analyses of the four trait groups, estimating simultaneously the phylogenetic signal for all traits and the correlations between traits, revealed a statistically significant phylogenetic signal for two of the four groups (photosynthesis and plant growth), comprising seven traits in total.

5. Our results demonstrate that even when the number of species in a comparative study is small, resulting in low power for univariate tests, phylogenetic signal can nonetheless be detected with multivariate tests that incorporate measurement error. Furthermore, our simulations show that the joint estimation of phylogenetic signal and trait correlations can lead to better (less biased and more precise) estimates of both.

Key-words: character syndromes, comparative methods, Magnoliaceae, phylogenetic inertia, phylogenetic signal, shade tolerance, strategy

*Correspondence authors. E-mail: arives@wisc.edu, caok@xtbg.cn

Introduction

Statistical methods that incorporate phylogenetic information are now common in comparative analyses of trait variation and covariation (e.g. Clobert, Garland & Barbault 1998; Housworth, Martins & Lynch 2004; Hansen, Pienaar & Orzack 2008; Lavin *et al.* 2008), because these methods enhance both biological insight and statistical validity. Biologically, shared ancestry should cause related species to resemble each other for many traits, a pattern referred to as phylogenetic signal (Blomberg & Garland 2002). For example, in a survey of traits taken from many studies, Blomberg, Garland & Ives (2003) demonstrated that phylogenetic signal was pervasive. Nonetheless, they found that phylogenetic signal was weaker on average for behavioural traits than for body size or size-adjusted morphometric traits. This finding may indicate that behavioural traits experience selection that breaks down phylogenetic patterns (Revell, Harmon & Collar 2008), giving inferential information about evolution of behavioural traits (although the possibility that they exhibit greater measurement error cannot be discounted; Ives, Midford & Garland 2007).

Statistically, analyses that require interspecific comparisons are challenging, because species cannot be assumed to be independent data points, violating the foremost assumption of many standard statistical analyses. When standard statistical methods are applied to phylogenetically related data, type I errors (rejecting the null hypothesis when in fact it is true) are often inflated, and coefficients estimated from statistical models (such as regression slopes) may not be minimum variance (e.g. Grafen 1989; Diaz-Uriarte & Garland 1996; Garland & Diaz-Uriarte 1999; Rohlf 2006). To address both biological and statistical issues, a large number of phylogenetically informed approaches have been developed for a wide range of analyses (reviewed in Garland, Bennett & Rezende 2005; Lavin *et al.* 2008).

Here, we extend this programme by developing new methods that estimate the correlation in trait values among species while simultaneously estimating the strength of phylogenetic signal and accounting for measurement error. This overcomes a limitation of the most commonly used phylogenetic statistical methods; they require an *a priori* assumption about the strength of phylogenetic signal in the data to be analysed. For example, Felsenstein's independent contrasts method (Felsenstein 1985) assumes that evolution follows a 'Brownian motion' process in which trait values increase or decrease randomly as evolution proceeds incrementally up a hierarchical phylogenetic tree. The assumption would be invalid, however, if trait evolution did not proceed in a Brownian motion fashion (e.g. Diaz-Uriarte & Garland 1996). In contrast, our methods estimate phylogenetic signal simultaneously with trait correlations, rather than making *a priori* assumptions about its strength, and so they should improve the estimation of trait correlations (Ackerly 1999) and related inferences, such as determining whether correlations differ statistically from zero.

Our methods are related to recently developed approaches used for detecting phylogenetic signal (Hansen 1997; Ackerly & Reich 1999; Freckleton, Harvey & Pagel 2002; Blomberg, Garland & Ives 2003; Butler & King 2004, 2005; Hansen, Pienaar & Orzack 2008; Scales, King & Butler 2009) and for performing regression analyses while simultaneously estimating the strength of phylogenetic signal (Lavin *et al.* 2008). Our methods use a conceptually related approach, but applied to correlation analyses. The approach is similar to that proposed by Freckleton, Harvey & Pagel (2002) and generalized by Revell & Harrison (2008), although it is based on an explicit model of evolution (see Blomberg, Garland & Ives 2003).

In addition to eliminating the need to make *a priori* assumptions about the strength of phylogenetic signal, our methods have two advantages. First, they allow for the joint estimation of phylogenetic signal in multiple traits (as does Freckleton, Harvey & Pagel 2002). Based on simulations of single continuous-valued traits, Blomberg, Garland & Ives (2003) found that *c.* 20 species are required to detect phylogenetic signal with a statistical power of about 0.8. However, statistical power could be improved if several traits are analysed simultaneously. For example, phylogeny aside, a multivariate analysis of variance (MANOVA) of several (correlated) traits typically yields much higher power to detect group differences compared with an ANOVA of a single trait (Willig & Owen 1987; Schmitz, Cherny & Fulker 1998). Our statistical approach allows for the joint estimation of phylogenetic signal for any group of traits, but rather than create groups arbitrarily, it makes most sense to combine traits based on level of biological organization (e.g. behaviour or biochemistry), functional relations (e.g. traits involved in locomotor or feeding performance), developmental origins (e.g. Dohm & Garland 1993) or genetic underpinnings (e.g. Arnold *et al.* 2008), because traits within such groups are expected to be correlated.

Second, our methods make it possible to incorporate within-species variation (or measurement error *sensu* Ives, Midford & Garland 2007). Real data sets contain within-species variation, and Ives, Midford & Garland (2007) and Felsenstein (2008) have shown that incorporating measurement error improves the accuracy of statistical tests and increases the statistical power to detect phylogenetic signal, group differences and so forth. By incorporating independent estimates of measurement error, our methods should give better estimates of trait correlations and phylogenetic signal.

We illustrate these methods by analysis of 13 traits measured in a common-garden study of nine species of *Manglietia* (Magnoliaceae). Below, we first focus on phylogenetic signal, asking which of 13 physiological and morphological traits show interspecific variation that to some extent reflects the relationships of the underlying phylogenetic tree. We then address the estimation of correlations among traits. For both phylogenetic signal and trait correlations, we present simulation studies to explore the statistical properties of the methods.

Materials and methods

PLANT TRAITS AND PHYLOGENETIC RECONSTRUCTION

We performed a common-garden experiment using nine *Manglietia* species: *M. grandis*, *M. hookeri*, *M. insignis*, *M. szechuanica*, *M. megaphylla*, *M. kwangtungensis*, *M. fordiana*, *M. chingii* and *M. pachyphylla* (for details, see Appendix S1, Supporting Information). After growing four seedlings of each species for 11 months in a greenhouse at the Xishuangbanna Tropical Botanical Garden, we performed 10 physiological and morphological measurements that divide into three categories: photosynthesis (maximum net photosynthesis A_{\max} , light saturation point LSP and light compensation point LCP), plant growth (basal stem diameter, crown volume, plant height, leaf area, relative growth rate) and thermal tolerance (critical temperature T_{ch} , peak temperature T_{\max} and temperature half-inactivation T_{50}). Data for the fourth group of traits, leaf morphology (thickness of leaves, palisade tissue, sponge tissue), were obtained for each species from the literature (Xie & Zhenghai 2000). Because all traits except the measurements of leaf morphology obtained from the literature were measured on multiple individuals, we used the standard error of the mean values of the traits for each species to estimate within-species variability, or measurement error (Ives, Midford & Garland 2007). Because multiple trait values were obtained from the same individuals, within-species variability could be correlated among different traits; for example, one individual that has a thick palisade layer might also have a thin sponge layer relative to other individuals in the same species. Therefore, we calculated the correlations in within-species variability among individuals and incorporated these into the statistical models along with the within-species variability.

Inferring phylogenetic relationships among closely related plant species is often difficult due to the lack of molecular markers exhibiting enough nucleotide variability. Therefore, we used many amplified fragment length polymorphisms (AFLPs) distributed throughout the whole genome, which proved capable of generating a hypothesis for the phylogenetic relationships among species (Appendix S1). For AFLP analyses, individual plants for each of the nine species ($n = 2-11$ individuals per species) were collected in the field or obtained from two botanical gardens, located in Kunming and Wenshan, Yunnan, China. We also included *Liriodendron chinense* ($n = 3$ individuals) as an outgroup to root the tree. A phylogenetic tree was constructed from the AFLP data using maximum likelihood with PAUP 4.0 (Swofford 1999). The analysis did not assume a molecular clock and resulted in a single maximum likelihood tree for the nine *Manglietia* species where the root position was determined by the location of the outgroup *Liriodendron*. The phylogenetic tree for the nine *Manglietia* species obtained using AFLP markers is presented in Fig. 1, and the AFLP data and the measurements of the 13 traits analysed in this article are given in Appendices S2 and S3.

PERMUTATION TEST FOR PHYLOGENETIC SIGNAL

A simple permutation test for phylogenetic signal is given by Blomberg & Garland (2002) and Blomberg, Garland & Ives (2003), and here we modify it to test for phylogenetic signal in multiple traits simultaneously. The permutation test is based on the null hypothesis that phylogenetic signal is absent, so that under the null hypothesis trait values can be permuted among species without changing the statistical characteristics of the data. For each permutation data set, the mean squared error (MSE) is calculated under the assumption




Fig. 1. Hypothesized phylogenetic tree for nine *Manglietia* species based on AFLP markers. Species codes are Mg, *M. grandis*; Mh, *M. hookeri*; Mf, *M. fordiana*; Mc, *M. chingii*; Mp, *M. pachyphylla*; Mi, *M. insignis*; Ms, *M. szechuanica*; Mm, *M. megaphylla*; and Mk, *M. kwangtungensis*. The tree is given in Nexus format with branch lengths in Appendix S2.

that phylogenetic signal exists. Specifically, the MSE is calculated assuming that evolution proceeds as a Brownian motion process using the specified hierarchical phylogenetic tree, so that the covariance in trait values between species is proportional to the amount of their shared phylogenetic ancestry (i.e. the branch-length distance from the root of the tree to their last common ancestor); more closely related species have a longer shared ancestry and hence greater predicted covariances in trait values. (The MSE calculated in this way is equivalent to the variance of standardized phylogenetically independent contrasts; Blomberg & Garland 2002; Blomberg, Garland & Ives 2003.) The distribution of MSE values calculated for the permutation data sets is compared with the MSE calculated for the observed data. If phylogenetic signal exists, then the MSE of the observed data will be distinctly low relative to the distribution of permutation MSEs, because incorporating phylogenetic structure (i.e. the assumption of Brownian motion evolution) leads to a better explanation (lower MSE) of the observed data.

Although Blomberg, Garland & Ives (2003) apply this test only to single traits taken separately, it can also be performed on multiple traits using the joint MSE for all traits. In this test, we want to weight each trait equally, so we standardized trait values x to have mean zero and variance 1, $z = \frac{x - \bar{x}}{\text{var}\{x\}^{1/2}}$. In other applications researchers might want to transform some traits, such as log-transform body size, in which case this should be done on the values of x before they are standardized.

PHYLOGENETIC SIGNAL AND TRAIT CORRELATIONS

Here, we describe the estimation of trait correlations with the simultaneous estimation of phylogenetic signal. The utility of this approach is that it leads to a full statistical model of both correlations among traits and phylogenetic correlations among species that can be used to test multiple hypotheses about trait correlations and phylogenetic signal. All of our computations for joint trait analyses were performed by the program CORRELATIONv2.m written in Matlab (Mathworks, Inc. 1996), available on request from T.G.

Our approach for the joint estimation of phylogenetic signal and trait correlations is based on a specific model of evolution under an Ornstein–Uhlenbeck (OU) process (Uhlenbeck & Ornstein 1930; Felsenstein 1988; Hansen 1997; Martins & Hansen 1997; Butler & King 2004; Scales, King & Butler 2009), although other models of evolution could be used (Grafen 1989; Martins & Hansen 1997; Pagel 1997;

Freckleton, Harvey & Pagel 2002; Housworth, Martins & Lynch 2004); for a review, see Lavin *et al.* (2008 Appendix A) and Revell, Harmon & Collar (2008). Our OU model is a multi-trait extension of the single-trait OU process as modelled by Blomberg, Garland & Ives (2003) in which evolution of the traits is correlated. This formulation differs slightly from that proposed by Martins & Hansen (1997), which assumes that the basal trait values are selected from the stationary distribution of an OU process. Instead, we assume that the variances in trait values at the base of the tree are zero, which has the advantage that our model becomes a Brownian motion model of evolution (which is a non-stationary process) as a special case.

To illustrate this model for two traits, suppose that over some arbitrarily small time step Δt , the evolution of trait values $x(t)$ and $y(t)$ for a given species is described by

$$\begin{aligned} X(t + \Delta t) &= d_x X(t) + \gamma_x(t) \\ Y(t + \Delta t) &= d_y Y(t) + \gamma_y(t) \end{aligned} \quad \text{eqn 1}$$

where d_x and d_y measure the strength of stabilizing selection for traits x and y , and $\gamma_x(t)$ and $\gamma_y(t)$ are random variables with means zero, variances σ_x^2 and σ_y^2 , and correlation r . Mapping this model onto a phylogenetic tree (under the assumption that the variance in trait values at the base of the tree is zero), the covariance in trait x between species i and j is

$$\text{cov}\{X_i, X_j\} = d_x^{\tau_i + \tau_j} \frac{1 - d_x^{2\tau_{ij}}}{1 - d_x^2} \sigma_x^2 \quad \text{eqn 2}$$

where X_i and X_j are the values of trait x for species i and j , τ_i and τ_j are the distances from the node representing their most recent common ancestor to the tip of the tree for species i and j , and τ_{ij} is the branch length from the base to the most recent common ancestor (Blomberg, Garland & Ives 2003). An identical expression holds for the covariance in values of trait y among species. Using a similar approach, it can be shown that the covariance between trait x for species i and trait y for species j is given by

$$\text{cov}\{X_i, Y_j\} = d_x^{\tau_i} d_y^{\tau_j} \frac{1 - (d_x d_y)^{\tau_{ij}}}{1 - d_x d_y} r \sigma_x \sigma_y \quad \text{eqn 3}$$

Thus, from the model of trait evolution (eqn 1) and a phylogenetic tree with N species, we can construct the variances and covariances in trait values among all species that incorporate both correlations between traits r and the strength of phylogenetic signal d . The stronger the stabilizing selection towards an optimum for a trait (the larger the phylogenetic signal parameter d), the weaker the phylogenetic correlation between species for that trait, because phylogenetic history is erased by selection towards the optimum (Felsenstein 1985; Blomberg, Garland & Ives 2003); a value of $d = 0$ corresponds to no phylogenetic signal (trait values among species are independent), and a value of $d = 1$ corresponds to Brownian motion evolution.

For use in statistical analyses, these variances (eqn 2) and covariances (eqn 3) must be combined into a covariance matrix. For illustrative purposes, consider hypothetical example of three species and two traits (Fig. 2). This leads to a 3×3 matrix \mathbf{C} whose diagonal elements τ_i contain the distance from the base to the tip for species i , and whose off-diagonal elements τ_{ij} give the shared distance on the phylogenetic tree between species i and j . In our example, we have standardized the branch lengths so that $\tau_i = 1$ (although the methods do not require that phylogenetic tree be ultra-metric, that is, have contempo-




Fig. 2. Construction of the joint trait-phylogeny covariance matrix that incorporates correlations among traits and correlations among phylogenetically related species. (a) The base information provided for a three-species data set with two traits x and y . The covariance matrix \mathbf{C} describes the phylogeny; the covariance in trait values between species is proportional to the shared branch lengths. (b) Strengths of phylogenetic signal when the values of $d_x = 2$ and $d_y = 0.25$ are estimated from the data. (Note that these values of d are hypothetical and in fact values of d cannot be estimated for data sets with fewer than four species.) The transformed matrices \mathbf{C}_x and \mathbf{C}_y incorporate the strengths of phylogeny signal which are depicted by the phylogenetic trees with transformed branch lengths. (c) The overall trait-species covariance matrix, $\sigma^2 \Psi$. The 3×3 submatrices along the diagonal of $\sigma^2 \Psi$ (shaded) give the covariance matrices containing correlations in values of a single trait among species. The 3×3 submatrices on the off-diagonal of $\sigma^2 \Psi$ give the covariance matrices containing correlations between traits x and y . The terms σ_x^2 and σ_y^2 give the variances in traits x and y , and r is the correlation between them.

aneous tips), and the shared branch between species 1 and 2 has length $\tau_{ij} = 0.7$ (Fig. 2a). In the statistical analysis, both the trait correlations r and the strength of phylogenetic signal d are estimated; for our example, there is a single value of r and two values of d , one for each trait. The effect of d on the strength of phylogenetic correlations can be visualized using a phylogenetic tree, but now with branch lengths proportional to the estimated phylogenetic signal. The value of phylogenetic signal for trait x , d_x , is 2, leading to a covariance of 0.8 between species, while for trait y , $d_y = 0.25$ and the covariance between species is 0.4; these covariances are contained within the covariance matrices $\mathbf{C}_x(d_x)$ and $\mathbf{C}_y(d_y)$ that are visualized by either extending (trait x) or contracting (trait y) the shared branch lengths between species (Fig. 2b). With $d_x = 2$ and $d_y = 0.25$, the correlation between traits is $r = 0.5$ and the variances for traits 1 and 2 are $\sigma_x^2 = 1.4$ and $\sigma_y^2 = 18.9$, resulting in the joint covariance matrix $\sigma^2 \Psi$ (Fig. 2c). Here, we use the standard convention of scaling the covariance matrix by σ^2 that gives the overall variance of the data (e.g. Judge *et al.* 1985).

The two blocks of the covariance matrix $\sigma^2 \Psi$ along the diagonal (shaded) correspond to the covariance matrices for traits x and y ; each matrix gives the variances for each of the three species along the diagonal and the phylogenetic covariances in the off-diagonals. The two blocks of matrix $\sigma^2 \Psi$ on the off-diagonals (not shaded) give the covariances among species between traits x and y . For example, the value 2.7 is the covariance between traits x and y expressed by species

1, while the value 1.7 is the covariance between trait x for species 1 and trait y for species 2. Thus, the matrix $\sigma^2\Psi$ gives covariances between all trait–species combinations.

The statistical method we developed estimates the components of the joint covariance matrix, specifically the parameters r , d and σ^2 , using Restricted Maximum Likelihood estimation; the details are presented in Appendix S4. The covariance matrix $\sigma^2\Psi$ can also be modified to incorporate measurement error. We treat as measurement error any within-species variation that might include error in the measurement of the trait or variation among individuals of the same species. (Population differences would also contribute to this error if they had been pooled to obtain a single sample to represent a species.) This measurement error is captured by the standard error of the within-species estimates of trait values (Ives, Midford & Garland 2007). To determine whether the estimates of d differ from zero or one, we used likelihood ratio tests. Specifically, we calculated the maximum restricted log likelihoods both including (LL) and excluding (LL₀) the parameters of interest, and tested the significance of the inclusion of the parameters using the result that $-2(LL_0 - LL)$ is asymptotically χ^2 -distributed with degrees of freedom equal to the number of parameters differing between models; for example, the test for $d \neq 0$ for three traits involves comparing the models with and without the three parameters d , and the resulting chi-squared distribution for $-2(LL_0 - LL)$ has 3 degrees of freedom.

In addition to estimating trait and phylogenetic correlations simultaneously for groups of traits, giving estimates we refer to as r_{joint} and d_{joint} , we modified the methods to calculate the value of d assuming that all species within the group have the same value d_{common} and estimate the corresponding trait correlations r_{common} . This provides a test for dissimilarity in phylogenetic signal among traits by comparing the fits of the model using d_{joint} and d_{common} . To compare with the jointly estimated values d_{joint} , we also computed the values of d estimated for each trait separately, d_{sep} ; we do not calculate the trait correlations r along with d_{sep} , because the correlations are components of the covariance matrix $\sigma^2\Psi$ for which values of d for each species should be estimated simultaneously (i.e. d_{joint}).

STATISTICAL POWER TO IDENTIFY PHYLOGENETIC SIGNAL

To compare the statistical power of the two methods (joint permutation test, likelihood ratio test) to detect phylogenetic signal in multiple traits, we performed a simulation study for three traits using the phylogenetic tree of the nine *Manglietia* species (Fig. 1). We simulated data sets under the assumption that evolution of the three traits followed an OU process with the same ‘true’ value of d for all traits. We assumed traits were not correlated ($r = 0$), although simulations with $r \neq 0$ gave similar results (data not presented). For each value of d between 0 and 1 in increments of 0.1, we simulated 2000 data sets, and for each simulated data set we performed the joint permutation test and the likelihood ratio test (obtained from the estimation of d_{joint}); we used an $\alpha = 0.05$ level to identify statistically significant phylogenetic signal.

STATISTICAL PROPERTIES OF ESTIMATORS OF D AND R

We investigated the statistical properties of the estimators of phylogenetic signal, d , and trait correlations, r , using simulations assuming that there are three traits under study for nine species with the same phylogeny as the nine *Manglietia* species (Fig. 1). To address the issue

of sample size, we also considered the case of 49 species using the phylogenetic tree provided by Garland *et al.* (1993). For both simulations, we assumed values of d are 0.2, 1 and 1.5, and values of r are 0.5, -0.5 and -0.25 . Finally, we simulated data both without and with measurement error; measurement error for the nine-species simulations were the same as for the nine *Manglietia* species, while for the 49-species simulations we assumed that the measurement error for each trait and the covariances between measurement errors were the same for all species and equal to the mean measurement errors for the corresponding traits in the nine *Manglietia* species data set. For the simulations with measurement error, we estimated d_{joint} and d_{sep} both incorporating and not incorporating measurement error.

Results

PHYLOGENETIC SIGNAL

The four different groups of traits (photosynthesis, leaf morphology, plant growth and thermal tolerance) demonstrated contrasting patterns of phylogenetic signal. The photosynthesis traits showed very strong phylogenetic signal (Table 1). For two of the three traits (A_{max} and LSP), the single-trait permutation tests were statistically significant and the estimates of d_{sep} for each trait analysed separately were greater than 1. The estimates of d_{joint} gave a similar pattern, and the joint permutation test was statistically significant. Finally, under the assumption that all three traits have the same value of d , the estimate of d_{common} was greater than 1 both with and without the inclusion of measurement error.

We used likelihood ratios to test three null hypotheses regarding the estimation of d_{joint} (Table 2): (i) whether there is phylogenetic signal in the group of traits (H_0 :no signal), (ii) whether the species covariances differ from those anticipated under Brownian motion evolution (H_0 :Brownian), and (iii) whether there is heterogeneity in values of d_{joint} among species (H_0 : $d_{\text{joint}} = d_{\text{common}}$). For the three photosynthesis traits, only the first of these hypotheses was rejected. We conclude that the phylogenetic signal is statistically indistinguishable from Brownian motion evolution, and that all three traits show indistinguishable degrees of phylogenetic signal.

For the three leaf morphology traits, the estimated values of d_{sep} and d_{joint} are large, yet they nonetheless cannot be statistically distinguished from zero; the separate likelihood ratio tests on d_{sep} (Table 1) and the joint likelihood ratio test for d_{joint} (Table 2) are not statistically significant. Nonetheless, the joint permutation test showed statistically significant phylogenetic signal (Table 1, $P < 0.03$). This illustrates that the tests using d and the permutation test do not necessarily give the same results for any given data set (see also Blomberg, Garland & Ives 2003). Below we demonstrate that the joint permutation test has greater statistical power than the likelihood ratio test for d_{joint} . Thus, it is not surprising that the permutation test identifies statistically significant phylogenetic signal while the likelihood ratio test does not. Despite the fact that the null hypothesis of no phylogenetic signal was rejected by the permutation test, we nonetheless rely solely

Table 1. Measures of phylogenetic signal for 13 plant traits. The permutation test gives a nonparametric assessment (P -value) for statistical departure from the case in which species are phylogenetically independent. Estimates of phylogenetic signal are given by the parameter d of an Ornstein–Uhlenbeck model of stabilizing selection, which is estimated for traits separately, d_{sep} , simultaneously for all traits within a group, d_{joint} , and simultaneously for all traits in a group under the assumption that all traits share the same value, d_{common} . Likelihood ratio tests were used to test the significant departure of d_{sep} from zero ($\dagger P < 0.1$; $*P < 0.05$; $**P < 0.02$), using likelihoods LL_{sep} and LL_0 calculated from the model including and excluding d_{sep} . The estimates of leaf morphology traits were taken from the literature and do not have associated measurement errors, and therefore values of d were not estimated with measurement error (shown by dashes in the table)

Trait	Permutation test (P -values)		d without ME					d with ME				
	Single	Joint	d_{sep}	LL_{sep}	LL_0	d_{joint}	d_{common}	d_{sep}	LL_{sep}	LL_0	d_{joint}	d_{common}
Photosynthesis												
A_{max}	0.03	0.03	4.55**	-3.69	-6.44	2.30	1.44	4.44†	-4.90	-6.67	1.82	1.67
LSP	0.02		2.66*	-38.29	-40.55	2.34		3.22*	-43.12	-45.25	3.93	
LCP	0.52		0.16	-8.54	-8.56	0.12		0.12	-9.59	-9.59	0.01	
Leaf morphology												
Thickness of leaves	0.21	0.04	2.65	-32.65	-33.01	0.74	0.26	–	–	–	–	–
Thickness of palisade tissue	0.14		1.98	-31.04	-31.45	1.36		–	–	–	–	–
Thickness of sponge tissue	0.14		1.85	-34.00	-34.30	0.23		–	–	–	–	–
Plant growth												
Basal stem diameter	0.36	0.59	0.24	-0.47	-0.48	0.14	1.52	0.47	-0.95	-0.97	1.51	0.83
Crown volume	0.66		0.11	-86.89	-86.89	1.39		0.11	-96.80	-96.80	0	
Leaf area	0.86		0.10	-48.29	-48.29	1.95		0.00	-52.92	-52.92	0.01	
Relative growth rate	0.28		0.44	-3.15	-3.24	1.97		0.46	-3.31	-3.37	0	
Thermal tolerance												
T_{ch}	0.74	0.88	0.08	-7.32	-7.32	0	0.00	0.00	-7.48	-7.48	0.30	0
T_{max}	0.96		0.41	-3.59	-3.59	0		0.00	-3.26	-3.26	0.04	
T_{50}	0.70		0.05	-5.25	-5.25	0.10		0.00	-5.25	-5.25	0.16	

Table 2. Joint tests for statistical significance of phylogenetic signal for four categories of traits. H_0 :no signal is the null hypothesis that there is no phylogenetic signal ($d = 0$ for all traits). H_0 :Brownian is the null hypothesis that the phylogenetic signal can be described by a Brownian motion model of evolution ($d = 1$ for all traits). H_0 : $d = d_{\text{common}}$ is the null hypothesis that the strength of phylogenetic signal is the same for all traits within a category. ME denotes measurement error and only P -values < 0.10 are shown. Degrees of freedom (d.f.) for the chi-squared tests equal the number of parameters that differ between models; for H_0 :no signal and H_0 :Brownian, d.f. = the number of traits per group, and for H_0 : $d = d_{\text{common}}$, d.f. = the number of traits - 1. Leaf morphology traits do not have associated measurement errors, and therefore models incorporating measurement error were not used

Trait category	Without ME					With ME						
	H_0 :no signal		H_0 :Brownian		H_0 : $d = d_{\text{common}}$	H_0 :no signal		H_0 :Brownian		H_0 : $d = d_{\text{common}}$		
	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P	χ^2	P
Photosynthesis	11.2	0.01	5.8		5.3		12.7	0.005	8.1	0.04	5.3	
Leaf morphology	2.34		2.57		2.17		–		–		–	
Plant growth	12.6	0.01	13.5	0.009	8.3	0.08	15.3	0.004	14.3	0.006	9.0	0.06
Thermal tolerance	0.14		6.9	0.015	0.14		3.3		6.0		3.3	

on the likelihood ratio test for reasons described below, concluding that there is no phylogenetic signal in leaf morphology traits.

The four traits for plant growth give an example in which the analyses of single traits revealed little evidence of phylogenetic signal, yet phylogenetic signal was found for the group of traits analysed jointly. Specifically, the null hypothesis H_0 :no signal was rejected whether measurement error was not ($P < 0.01$) or was ($P < 0.004$, Table 2) included. Moreover, the null hypothesis H_0 :Brownian is also rejected ($P < 0.009$ and < 0.006 with and without measurement error), indicating

that even though there is evidence for phylogenetic signal, the Brown motion evolutionary model is also not supported. The nearly significant rejection of the null hypothesis H_0 : $d = d_{\text{common}}$ ($P < 0.08$ and < 0.06 with and without measurement error) suggests that traits differ in whether or not they show phylogenetic signal, which is supported by the very different estimates among species of both d_{sep} and d_{joint} (Table 1). Note that in contrast to the leaf morphology traits, although the likelihood ratio tests identify joint phylogenetic signal, the permutation test does not. We return to this in the Discussion.

For the fourth category of traits (thermal tolerance), there is little indication of phylogenetic signal in any analysis.

STATISTICAL POWER TO IDENTIFY PHYLOGENETIC SIGNAL

In our simulation study, the joint permutation test had greater statistical power than the likelihood ratio test (Fig. 3), showing greater ability to reject the null hypothesis of no phylogenetic signal than the likelihood ratio test for $d > 0$. Furthermore, the likelihood ratio test had a slightly inflated type I error rate (probability of rejecting the null hypothesis when it is true), with 6.6% of the data sets rejected at the $\alpha = 0.05$ level when $d = 0$; this compared to a type I error rate of 5.6% for the permutation test. Importantly, only 2.0% of the simulated data sets were rejected by both the joint permutation test and the likelihood ratio test. Therefore, if one were to apply both tests to the same data set and conclude that the data set had phylogenetic signal if either one or the other of the tests rejected the null hypothesis, then the type I error rate would be 10.2%; the type I error rates for the permutation and likelihood ratio tests were 5.6% and 6.6%, so the probability of one or the other test rejecting the null hypothesis is $5.6\% + 6.6\% - 2.0\% = 10.2\%$. This suggests that it is necessary to make an *a priori* decision about which statistical test to use – and use only that one – in order to obtain the correct type I error rates.

TRAIT CORRELATIONS

We computed trait correlations within each of the four groups. Table 3 gives four types of correlation coefficients: (i) Pearson correlation coefficients obtained under the assumption that there is no phylogenetic signal; (ii) coefficients calculated under the assumption of Brownian motion evolution;




Fig. 3. Power analysis for the joint permutation and likelihood ratio tests for three traits simulated using the phylogenetic tree of nine *Manglietia* species (Fig. 1). Each trait was simulated using an Ornstein–Uhlenbeck evolutionary process with a common value of d , and traits were assumed to be independent. For values of d between zero and one in increments of 0.1, 2000 simulations were performed, and the figure gives the fraction for which the null hypothesis of $d = 0$ was rejected using the joint permutation test and likelihood ratio test at an $\alpha = 0.05$ level.

(iii) coefficients calculated at the same time as d_{joint} for all three traits, r_{joint} ; and (iv) r_{joint} calculated while incorporating measurement error. For brevity, we do not report correlations r_{common} obtained when assuming all species have a common value of d .

For all groups of traits, the values of the correlation coefficients computed using all four methods exhibit general correspondence (Table 3); this empirical correspondence between standard and phylogenetic correlations has been shown previously (Ricklefs & Starck 1996; Ackerly & Reich 1999). Furthermore, there was statistically significant correlation between traits within all four groups; all likelihood ratio tests of the hypothesis that all correlations are zero were rejected using all four of the approaches for measuring correlation coefficients (Table 3). Comparing statistical tests among the four approaches used for estimating correlation coefficients, the most striking result is the stronger rejections (lower P -values) of the null hypotheses when the analyses incorporated measurement error; for the three groups for which measurement error was available (photosynthesis, leaf morphology and thermal tolerance), the null hypotheses of zero correlations were rejected at the $P < 0.03$, 0.01 and 0.001 levels using Pearson correlations (assuming no measurement error and no phylogenetic signal), whereas all these hypotheses were rejected at the $P < 0.001$ level when estimating r_{joint} with measurement error (Table 3).

STATISTICAL PROPERTIES OF d AND r

To explore the statistical properties of the joint estimation of d and r , we performed a simulation study using phylogenetic trees for nine species (Fig. 1) and 49 species (Garland *et al.* 1993). For simulations without measurement error, the estimates of d_{joint} were less biased and more precise (i.e. less variable) than the estimates calculated for each species separately, d_{sep} , when there were nine species (Fig. 4a). Increasing the sample size to 49 removed any bias for both methods, and the estimates of d_{joint} are only very slightly more precise (Fig. 4b). Thus, for small sample sizes the estimates of d_{joint} are preferred over d_{sep} , even without the added advantage of d_{joint} that it provides statistical tests of phylogenetic signal for groups of traits. (We should of course note that this is true only for the specific simulations with the specific trait correlations r that we investigated, and that researchers might want to perform simulations based on their own data to confirm this. Nonetheless, we suspect this pattern will hold broadly.) In simulations including measurement error (Fig. 4c,d), the estimates of d that ignored the simulated measurement error were more biased than those incorporating measurement error, especially for the case with 49 species (Fig. 4d). Furthermore, the estimates of d_{joint} were more precise than those for d_{sep} for small sample sizes (Fig. 4c). Thus, for data sets with substantial measurement error, estimating d_{joint} with measurement error incorporated has the best statistical properties.

In the same simulations we used to estimate values of d (Fig. 4), we also obtained estimates of r (Fig. 5); we did not

	P-value	Trait pairs					
		1-2	1-3	2-3	1-4	2-4	3-4
(a) Photosynthesis traits							
No signal	0.04	0.63	-0.63	-0.30			
Brownian	0.005	0.71	-0.49	-0.18			
r_{joint}	0.03	0.62	-0.56	-0.26			
r_{joint} with ME	0.001	0.71	-0.52	0			
Ave. d_{joint} with ME		2.82	0.61	1.97			
Ave. d_{joint}		2.32	1.21	1.23			
(b) Leaf morphology traits							
No signal	0.002	0.34	0.61	-0.42			
Brownian	0.002	0.30	0.56	-0.47			
r_{joint}	0.003	0.25	0.52	-0.55			
Ave. d_{joint}		1.04	0.47	0.80			
(c) Growth traits							
No signal	0.01	0.30	0.29	0.37	0.84	0.65	0.75
Brownian	0.01	0.32	0.24	0.40	0.92	0.60	0.74
r_{joint}	0.001	0.35	0.29	0.92	0.45	0.63	0.83
r_{joint} with ME	0.001	0.26	-0.02	0.36	0.32	0.83	0.54
Ave. d_{joint} with ME		0.75	0.76	0.01	0.75	0.00	0.01
Ave. d_{joint}		0.77	1.05	1.67	1.06	1.68	1.96
(d) Thermal tolerance traits							
No signal	0.001	0.80	0.85	0.64			
Brownian	0.01	0.82	0.86	0.68			
r_{joint}	0.001	0.80	0.85	0.64			
r_{joint} with ME	0.001	0.50	0.99	0.52			
Ave. d_{joint}		0	0.05	0.05			
Ave. d_{joint} with ME		0.17	0.23	0.10			

Table 3. Correlation coefficients calculated using four methods. The 'no signal' and 'Brownian motion' calculations assume that there is no phylogenetic signal ($d = 0$) and signal given by Brownian motion evolution along the original specified phylogenetic tree ($d = 1$). Coefficients were also calculated using phylogenetic signal estimated simultaneously for all traits, d_{joint} , with and without incorporating measurement error. When d was estimated, the average of the estimates of d is provided for each pair of species. The traits are listed in order: (a) photosynthesis traits (A_{max} , LSP, LCP), (b) leaf morphology traits (thickness of leaves, thickness of palisade tissue, and thickness of sponge tissue), (c) growth traits (basal stem diameter, crown volume, leaf area, relative growth rate) and (d) thermal tolerance traits (critical temperature T_{ch} , peak temperature T_{max} , temperature half-inactivation T_{50}). P-values correspond to the likelihood ratio test for the null hypothesis $H_0: r = 0$ for all species pairs

include estimates of r when estimating d_{sep} for reasons described in Material and methods. In simulations not including measurement error (Fig. 5a,b), the estimates of r under the assumption of no phylogenetic signal or Brownian motion evolution were both biased for the larger sample size of 49 species, whereas the estimates of r_{joint} were not. In simulations with measurement error (Fig. 5c,d), with 49 species the estimates using all methods were biased except for the joint estimate of r_{joint} incorporating measurement error. These results illustrate the relatively good statistical performance of estimators of d_{joint} and r_{joint} that simultaneously include phylogenetic signal and trait correlations.

Discussion

Our statistical methods to measure phylogenetic signal in multiple traits simultaneously revealed statistically significant signal in the group of three traits involved in photosynthesis (A_{max} , LSP and LCP) and the group of four traits involving plant growth (basal stem diameter, crown volume, leaf area and relative growth rate). A third group of three traits involving leaf morphology showed phylogenetic signal in the permutation test but not in the likelihood ratio tests; estimates of measurement error were not available, so our method for incorporating measurement error could not be applied. Finally, the fourth group of thermal tolerance traits exhibited no phylogenetic signal. These results from analyses that consider traits jointly within the same group contrast the conclusions obtained when analysing the 13 traits separately. Analysed separately, we found strong phylogenetic signal in

only two traits, A_{max} and LSP, both of which involve photosynthesis. Failure to detect phylogenetic signal in traits treated separately is not surprising, however, because statistical tests of phylogenetic signal often lack power when sample sizes are small (Blomberg, Garland & Ives 2003).

For the groups of traits showing statistically significant phylogenetic signal, all we conclude is that phylogenetically related species are more likely to share the same values of the group of traits. Although our methods rely on a specific model of evolution, assuming trait values evolve according to an OU process of stabilizing selection across the phylogeny (eqn 1), it would be an over-interpretation of the results to conclude that the data were produced under an OU process. A formal statistical test of the specific model of evolution would consist of constructing competing models derived under different evolutionary processes and then selecting the best amongst them; although this procedure is straightforward, we suspect that large amounts of high-quality data would be required to statistically distinguish among models. Our goal, however, was only to identify the existence of phylogenetic signal, not to test any specific mechanisms that underlie phylogenetic patterns (cf. Blomberg & Garland 2002; Blomberg, Garland & Ives 2003).

Previous studies of plants have documented cases of both strong phylogenetic signal and weak or absent signal. Ackerly & Reich (1999) assembled a data set of 108 tree species for which eight leaf traits were available. All of the traits showed phylogenetic signal, although this was largely due to the data set spanning angiosperms and gymnosperms, two clades with distinct leaf morphology (broad leaves vs. needles); in similar




Fig. 4. Estimates of phylogenetic signal, d , for simulated data for three traits. In (a) and (b) data were simulated without measurement error, and d_{sep} and d_{joint} were estimated excluding measurement error. In (c) and (d) measurement error was simulated, and both d_{sep} and d_{joint} were estimated including and not including measurement error. In (a) and (c) the phylogenetic tree for the nine *Manglietia* species was used for simulations, and in (b) and (d) the tree for 49 species from Garland *et al.* (1993) was used. Points give the mean of the estimates from simulations, vertical lines give the 95% inclusion intervals and horizontal lines give the true values of d for the three traits used in the simulations: 0.2, 1 and 1.5. Values of $r = 0.5, -0.5$ and -0.25 . For (a) and (c) 1000 data sets were simulated, and for (b) and (d) 200 data sets were simulated.

analyses confined to angiosperms, they found little phylogenetic signal in the same traits. Zanne, Chapman & Kitajima (2005) found strong phylogenetic signal in cotyledon type but not seed mass among 70 species of trees and shrubs occupying the same habitat in Uganda. In a survey of pollination traits among 288 species, Ornelas *et al.* (2007) found phylogenetic signal in all traits involving nectar production, as well as corolla length, suggesting strong evolutionary conservatism in plant reproductive biology.

These studies were broad taxonomic surveys, however, involving diverse plant lineages. In contrast, our study compared a small group of congeners. At a similar taxonomic scale, Blomberg, Garland & Ives (2003) analysed two plant data sets consisting of morphological traits for maple trees (Ackerly & Donoghue 1998) and *Tithonia* (Asteraceae) (Morales 2000). Despite the larger size of these data sets (17 and 32 species vs. our 9), phylogenetic signal was found for only about one-third of traits (4 of 12 and 5 of 14 traits respectively) based on the same permutation test that we employed here (with P -values not corrected for multiple comparisons). In a common-garden experiment on drought tolerance of eight species from five genera, Valladares & Sanchez-Gomez (2006) documented phylogenetic signal in several traits (as




Fig. 5. Estimates of the correlation coefficients for simulated data sets for three traits. The same data sets were used as those producing corresponding panels in Fig. 4. In (a) and (b) data were simulated without measurement error, and d_{sep} and d_{joint} were estimated excluding measurement error. The correlation coefficients were then estimated assuming no phylogenetic signal, Brownian motion evolution and phylogenetic signals given by d_{joint} . In (c) and (d) measurement error was simulated, and d_{joint} was estimated including and not including measurement error. In (a) and (c) the phylogenetic tree for the nine *Manglietia* species was used for simulations, and in (b) and (d) the tree for 49 species from Garland *et al.* (1993) was used. Points give the mean of the estimates from simulations, vertical lines give the 95% inclusion intervals and horizontal lines give the true values of r for the three traits (0.5, -0.5 and -0.25). For (a) and (c) 1000 data sets were simulated, and for (b) and (d) 200 data sets were simulated.

determined by clustering of traits in a Principle Components Analysis), with similar drought tolerance exhibited by members of the same genera. In comparison with the taxonomically broad surveys, these studies suggest that phylogenetic signal in plant traits is harder to identify among taxonomically similar species. However, this may also be the result of sample size, as confining analyses to closely related species often limits the number of species that can be analysed, thus reducing statistical power.

Our statistical methods provide a means for estimating simultaneously the correlations of multiple traits and the phylogenetic correlations in values of traits among species. In a simulation study, we showed that correlations calculated when either ignoring phylogenetic signal or assuming phylogenetic signal generated a Brownian motion model of evolution often produced biased estimates of the true correlation (see also Rohlf 2006). In contrast, estimating trait correlations while simultaneously estimating phylogenetic signal, r_{joint} , gave less-biased estimates. This demonstrates that correctly accounting for phylogenetic signal, rather than making an *a priori* assumption that it is zero or given

by Brownian motion evolution, leads to better estimates of correlations.

Although we present two tests for phylogenetic signal in suites of traits, the joint permutation test and the likelihood ratio test using estimates of d_{joint} , only one of these tests should be selected *a priori* and used to determine statistical significance. This is because the two tests do not necessarily identify the same data sets as showing phylogenetic signal in suites of traits (see also Blomberg, Garland & Ives 2003). To give an extreme example, if both tests had rejection rates of the null hypothesis of 5%, yet data sets rejected by one test were never rejected by the other, then 10% of the data sets would be rejected by one or the other test. Therefore, if a researcher were to assume that a data set showed phylogenetic signal for a suite of traits if one or the other statistical test gave statistical significance, then in this example the type I error rate (the rate of rejecting a null hypothesis that is true) would be inflated to 10%. In our simulations, only 2.0% of simulated data sets had null hypotheses that were rejected by both tests, indicating potentially serious type I errors if both tests were applied. The underlying explanation for this statistical complexity is that the permutation and likelihood ratio tests are rejecting the null hypothesis based on different characteristics of the data sets; they are employing different alternative (H_1) hypotheses.

Given that either the permutation test or the likelihood ratio test should be selected *a priori* as the group test for phylogenetic signal, which one should it be? The joint permutation test has the advantage of greater power, and therefore it might be preferred if detecting phylogenetic signal is the only objective. The likelihood ratio test has the advantage of being a model-based test that gives more information about the data set, including an estimate of the strength of phylogenetic signal and phylogenetically correct correlations among traits. In our study, we were interested in not only phylogenetic signal but also trait correlations, and therefore we would pick the likelihood ratio test over the permutation test. Thus, although the permutation test shows phylogenetic signal for the group of leaf morphology traits, we do not conclude that they in fact exhibit phylogenetic signal.

The product of the estimation of d_{joint} and r_{joint} is the joint covariance matrix $\sigma^2\Psi$ that incorporates correlations due to phylogenetic relationships among species, associations between traits and measurement error (Fig. 1, Appendix S1). This covariance matrix can be best explained by considering each value of a trait for each species as a trait–species datum. The elements of the covariance matrix are then the covariances between each pair of trait–species values. In the structure assumed for the covariance matrix $\sigma^2\Psi$, trait–species values might be correlated if species are phylogenetically related. They might also be correlated if the two traits involved in the trait–species values have experienced correlated evolution along the phylogenetic tree. Although the structure we assume for the covariance matrix is flexible to include different strengths of phylogenetic signal and different trait correlations, it nonetheless imposes constraints on the covariances. For example, it assumes that closely related

species are at least as likely to share similar trait values as more distantly related species. This assumption might be violated, however, if there is convergent evolution, in which distantly related species have become similar in multiple traits. Although the structure of the covariance matrix $\sigma^2\Psi$ cannot accommodate convergent evolution, it can nonetheless be used to provide a null hypothesis that no convergence exists. This could be done, for example, by fitting the covariance matrix $\sigma^2\Psi$ to a data set, using the fitted model to simulate data, and then looking for patterns of convergence in the observed data that are not found in the simulated data; methods to do this, however, will require further development. In general, by fitting an explicit model to the covariance structure of a data set, our approach provides an avenue to investigate complex evolutionary hypotheses about the variation in multiple traits among phylogenetically related species.

Acknowledgements

We thank especially Stacey Smith for comments and help with this project. It was supported in part by NSF grants DEB-0816613 to ARI and DEB-0416085 to D.N. Reznick, M.S. Springer and T.G.

References

- Ackerly, D.D. (1999) Comparative plant ecology and the role of phylogenetic information. *Physiological Plant Ecology* (eds M. Press, J. Scholes & M. Barker), pp. 391–413. Blackwell Science, Oxford.
- Ackerly, D.D. & Donoghue, M.J. (1998) Leaf size, sapling allometry, and Corner's rules: phylogeny and correlated evolution in maples (*Acer*). *The American Naturalist*, **152**, 767–791.
- Ackerly, D.D. & Reich, P.B. (1999) Convergence and correlations among leaf size and function in seed plants: a comparative test using independent contrasts. *American Journal of Botany*, **86**, 1272–1281.
- Arnold, S.J., Burger, R., Hohenlohe, P.A., Ajie, B.C. & Jones, A.G. (2008) Understanding the evolution and stability of the G-matrix. *Evolution*, **62**, 2451–2461.
- Blomberg, S.P. & Garland, T., Jr (2002) Tempo and mode in evolution: phylogenetic inertia, adaptation and comparative methods. *Journal of Evolutionary Biology*, **15**, 899–910.
- Blomberg, S.P., Garland, T., Jr & Ives, A.R. (2003) Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution*, **57**, 717–745.
- Butler, M.A. & King, A.A. (2004) Phylogenetic comparative analysis: a modeling approach for adaptive evolution. *The American Naturalist*, **164**, 683–695.
- Butler, M.A. & King, A.A. (2005) Ouch! An improved method for phylogenetic tests of adaptive evolution. *Integrative and Comparative Biology*, **45**, 973–973.
- Clobert, J., Garland, T., Jr & Barbault, R. (1998) The evolution of demographic tactics in lizards: a test of some hypotheses concerning life history evolution. *Journal of Evolutionary Biology*, **11**, 329–364.
- Diaz-Uriarte, R. & Garland, T., Jr (1996) Testing hypotheses of correlated evolution using phylogenetically independent contrasts: sensitivity to deviations from Brownian motion. *Systematic Biology*, **45**, 27–47.
- Dohm, M.R. & Garland, T., Jr (1993) Quantitative genetics of scale counts in the garter snake *Thamnophis sirtalis*. *Copeia*, **4**, 987–1002.
- Felsenstein, J. (1985) Phylogenies and the comparative method. *The American Naturalist*, **125**, 1–15.
- Felsenstein, J. (1988) Phylogenies and quantitative characters. *Annual Review of Ecology and Systematics*, **19**, 445–471.
- Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002) Phylogenetic analysis and comparative data: a test and review of evidence. *The American Naturalist*, **160**, 712–726.
- Garland, T., Jr, Bennett, A.F. & Rezende, E.L. (2005) Phylogenetic approaches in comparative physiology. *Journal of Experimental Biology*, **208**, 3015–3035.

- Garland, T., Jr & Díaz-Uriarte, R. (1999) Polytomies and phylogenetically independent contrasts: an examination of the bounded degrees of freedom approach. *Systematic Biology*, **48**, 547–558.
- Garland, T., Jr, Dickerman, A.W., Janis, C.M. & Jones, J.A. (1993) Phylogenetic analysis of covariance by computer-simulation. *Systematic Biology*, **42**, 265–292.
- Grafen, A. (1989) The phylogenetic regression. *Transactions of the Royal Society of London B, Biological Sciences*, **326**, 119–157.
- Hansen, T.F. (1997) Stabilizing selection and the comparative analysis of adaptation. *Evolution*, **51**, 1341–1351.
- Hansen, T.F., Pienaar, J. & Orzack, S.H. (2008) A comparative method for studying adaptation to a randomly evolving environment. *Evolution*, **62**, 1965–1977.
- Housworth, E.A., Martins, E.P. & Lynch, M. (2004) The phylogenetic mixed model. *The American Naturalist*, **163**, 84–96.
- Ives, A.R., Midford, P.E. & Garland, T., Jr (2007) Within-species variation and measurement error in phylogenetic comparative methods. *Systematic Biology*, **56**, 252–270.
- Judge, G.G., Griffiths, W.E., Hill, R.C., Lutkepohl, H. & Lee, T.-C. (1985) *The Theory and Practice of Econometrics*. John Wiley and Sons, New York.
- Lavin, S.R., Karasov, W.H., Ives, A.R., Middleton, K.M. & Garland, T., Jr (2008) Morphometrics of the avian small intestine, compared with non-flying mammals: a phylogenetic approach. *Physiological and Biochemical Zoology*, **81**, 526–550.
- Martins, E.P. & Hansen, T.F. (1997) Phylogenies and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. *The American Naturalist* **149**, 646–667. Erratum 153:448.
- Mathworks, Inc. (1996) *MATLAB*. The MathWorks, Inc., Natick, MA.
- Morales, E. (2000) Estimating phylogenetic inertia in *Tithonia* (Asteraceae): a comparative approach. *Evolution*, **54**, 475–484.
- Ornelas, J.F., Ordano, M., De-Nova, A.J., Quintero, M.E. & Garland, T., Jr (2007) Phylogenetic analysis of interspecific variation in nectar of hummingbird-visited plants. *Journal of Evolutionary Biology*, **20**, 1904–1917.
- Pagel, M. (1997) Inferring evolutionary processes from phylogenies. *Zoologica Scripta*, **26**, 331–348.
- Revell, L.J., Harmon, L.J. & Collar, D.C. (2008) Phylogenetic signal, evolutionary process, and rate. *Systematic Biology*, **57**, 591–601.
- Revell, L.J. & Harrison, A.S. (2008) PCCA: a program for phylogenetic canonical correlation analysis. *Bioinformatics*, **24**, 1018–1020.
- Ricklefs, R.E. & Starck, J.M. (1996) Applications of phylogenetically independent contrasts: a mixed progress report. *Oikos*, **77**, 167–172.
- Rohlf, F.J. (2006) A comment on phylogenetic correction. *Evolution*, **60**, 1509–1515.
- Scales, J.A., King, A.A. & Butler, A.M. (2009) Running for your life or running for your dinner: what drives fiber-type evolution in lizard locomotor muscles? *The American Naturalist*, **173**, 543–553.
- Schmitz, S., Cherny, S.S. & Fulker, D.W. (1998) Increase in power through multivariate analyses. *Behavior Genetics*, **28**, 357–363.
- Swofford, D.L. (1999) *PAUP: Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Sinauer Associates, Inc., Sunderland, MA.
- Uhlenbeck, G.E. & Ornstein, L.S. (1930) On the theory of Brownian motion. *Physical Review*, **36**, 823–841.
- Valladares, F. & Sanchez-Gomez, D. (2006) Ecophysiological traits associated with drought in Mediterranean tree seedlings: Individual responses versus interspecific trends in eleven species. *Plant Biology* **8**, 688–697.
- Willig, M.R. & Owen, R.D. (1987) Univariate analyses of morphometric variation do not emulate the results of multivariate analyses. *Systematic Zoology*, **36**, 398–400.
- Xia, C. & Zhenghai, H. (2000) Comparative studies on leaf structure and oil cells of the Magnoliaeae in China. *Acta Phytotaxonomica Sinica*, **38**, 218–230.
- Zanne, A.E., Chapman, C.A. & Kitajima, K. (2005) Evolutionary and ecological correlates of early seedling morphology in East African trees and shrubs. *American Journal of Botany*, **92**, 972–978.

Received 20 January 2009; accepted 19 May 2009

Handling Editor: James Cresswell

Supporting information

Additional supporting information may be found in the online version of this article:

Appendix S1. Methodological details of the common-garden experiments and AFLP analysis for phylogenetic tree construction for nine *Manglietia* species.

Appendix S2. AFLP data for nine *Manglietia* species and the outgroup *Liriodendron chinense* used to construct the phylogenetic tree (Fig. 2), including the phylogenetic tree in Nexus form.

Appendix S3. Data for 13 traits for the nine *Manglietia* species.

Appendix S4. Derivations for the statistical methods used in the analyses.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

APPENDIX S1: METHODOLOGICAL DETAILS OF THE COMMON-GARDEN EXPERIMENTS AND AFLP ANALYSIS FOR PHYLOGENETIC TREE CONSTRUCTION FOR NINE *MANGLIETIA* SPECIES

COMMON-GARDEN EXPERIMENT

To measure trait values, we performed a common-garden experiment using nine *Manglietia* species: *M. grandis*, *M. hookeri*, *M. insignis*, *M. szechuanica*, *M. megaphylla*, *M. kwangtungensis*, *M. fordiana*, *M. chingii*, and *M. pachyphylla*. Seeds were collected at the Fadou Shan Natural Reserve in the Xiangpingshan forest of Xichou county, Yunnan, China in September, 2005, germinated on seedbeds, and then transplanted to pots in a mixture of forest surface soil and river sand in May, 2006. Transplanted seedlings, approximately 15 plants per species, were given slow-release fertilizer and grown under common-garden conditions in a greenhouse at the Xishuangbanna Tropical Botanical Garden, the Chinese Academy of Sciences, Yunnan, China. In the following spring, after 11 months of growth, we performed 10 physiological and morphological measurements that divide into three categories: photosynthesis (maximum net photosynthesis A_{max} , light saturation point LSP, and light compensation point LCP), plant growth (basal stem diameter, crown volume, plant height, leaf area, relative growth rate), and thermal tolerance (critical temperature T_{ch} , peak temperature T_{max} , and temperature half-inactivation T_{50}).

For measurements of photosynthesis traits and thermal tolerance (gas exchange experiments), one leaf per individual for 4 individuals per species was sampled within each plot. Gas exchange was measured midday (10:00 to 15:00) with a LI-COR 6400 portable photosynthesis system (LI-COR, Lincoln, Nebraska, USA). Air was supplied at a constant flow

rate, and CO₂ concentration was at the naturally occurring level. A built-in LED red light source was used to produce different irradiances during each measurement. Photosynthesis-irradiance curves were generated with photon flux densities (PFDs) of 1500, 1200, 1000, 800, 500, 300, 200, 100, 40, 20, and 0 mol/m² s. In each trial, a leaf was placed in the chamber and allowed to acclimate to the maximum light level, 2000 mol/m² s. Each subsequent measurement was delayed 2-6 min for equilibration before the light level was changed. Photosynthesis-irradiance curves were constructed using rates of gas exchange measured at the 11 PFDs by fitting a nonrectangular hyperbola, and from these curves the asymptotic light-saturated rate of net photosynthesis (A_{max}), near-light saturation point (LSP) and light compensation point (LCP) were determined. To measure thermal tolerance, the temperature of the chamber was then increased from 20 to 65 C. The critical temperature (T_{ch}), peak temperature (T_{max}), and temperature half-inactivation (T₅₀) for heat tolerance of photosystem 2 were determined graphically from the F₀ to leaf temperature plots (Braun, Buchner & Neuner, 2002). Temperature half-inactivation (T₅₀) is the temperature at which 50% of leaf tissue is heat damaged (Braun *et al.*, 2002).

Plant growth measurements (basal stem diameter, crown volume, leaf area, relative growth rate) were taken on the 11-month-old saplings. Basal stem diameter was measured at 1/10 of the stem height. Crown volume was estimated from an ellipsoid with dimensions D_1 , D_2 and D_3 , where D_1 and D_2 are the diameter of the crown in the major and minor axes, and D_3 is the depth of the foliage; crown volume equals $\pi D_1 D_2 D_3 / 6$. Total leaf area was estimated using a LI 3000A Area Meter (LI-COR, Lincoln, Nebraska, USA). The relative growth rate (RGR) was calculated as $RGR = \ln(W_{\text{final}} - \ln W_{\text{initial}}) / (T_{\text{final}} - T_{\text{initial}})$, where W_{final} and W_{initial} are dry biomass

at the time of harvest (T_{final}) and the time of transplanting (T_{initial}) from seedbed into pots (Hunt, 1982).

Data for the fourth group of traits, leaf morphology (thickness of leaves, palisade tissue, sponge tissue), were obtained for each species from the literature (Xia & Zhenghai, 2000).

PHYLOGENY RECONSTRUCTION

Inferring phylogenetic relationships among closely related plant species is often difficult due to the lack of molecular markers exhibiting enough nucleotide variability. A solution to this problem is to use many amplified fragment length polymorphisms (AFLPs) distributed throughout the whole genome. Althoff et. al. (2007) concluded that AFLP data are best-suited for examining phylogeographic patterns within species and among very recently diverged species. Our previous analyses (Zheng Li and Kunfang Cao, unpublished) found that noncoding chloroplast DNA sequences were not informative at the intrageneric level, and internal transcribed spacer (ITS) sequences could not be amplified. In contrast, AFLP proved capable of generating a hypothesis for the phylogenetic relationships among species.

For AFLP analyses, individual plants were collected in the field or obtained from two botanical gardens, located in Kunming and Wenshan, Yunnan, China. Individuals from the same species were collected from different locations to ensure little genetic relatedness among individuals within species. Sample sizes for the nine species were *M. grandis* ($n = 6$), *M. hookeri* ($n = 11$), *M. insignis* ($n = 10$), *M. szechuanica* ($n = 13$), *M. megaphylla* ($n = 2$), *M. kwangtungensis* ($n = 8$), *M. fordiana* ($n = 11$), *M. chingii* ($n = 2$), and *M. pachyphylla* ($n = 2$). We also included *Liriodendron chinense* ($n = 3$) as an out group to root the tree. Genomic DNA was extracted following the method of Doyle and Doyle (1987). This generated a genotyping set

of 65 individuals. All extracted genomic DNA was purified using Qiagen Quick PCR Purification Kit (Qiagen, Shanghai, China). Total genomic DNA was digested with two restriction enzymes (EcoRI and MseI). The AFLP procedure was performed as described in Vos et al. (1995) using preselective PCR primers E.A: 5'-GACTGCGTACCAATTCA-3' and M.C: 5'-GATGAGTCCTGAGTAAC-3'. Three selective PCR primer pairs were chosen from 32 tested primers based on the quality of the produced bands (i.e., even distribution of bands with relatively homogeneous intensity): E.AGC/M.CTT, E.AAG/M.CAC, and E.AAG /M.CTG. Amplified fragments were separated on a 6% denaturing polyacrylamide gel at 110 W using Sequi-Gen® GT Sequencing Cell (Bio-Rad, Hercules, California, USA), and bands were visualized by silver nitrate staining. Care was taken to ensure uniformity of band intensity between gels by stopping the developing reaction with reference to the selected standard bands within the profile. Positions of AFLP bands within a readable range of 50-500 bp were recorded as a binary character matrix (0/1 for presence/absence of a band at a particular position).

Maximum likelihood analyses were conducted with PAUP 4.0 (Swofford, 1999). For this analysis, 0/1 characters were coded as purines and pyrimidines with a single set of markers for each species. For inclusion as a character value of 1, an AFLP band had to be present in all individuals of the same species. The analysis did not assume a molecular clock and resulted in a single maximum likelihood tree for the nine *Manglietia* species where the root position was determined by the location of the outgroup *Liriodendron*. This maximum likelihood tree has the same tree topology as the majority rule consensus tree from an analysis of the same data using a two-state continuous-time Markov model in MrBayes version 3.1 (Huelsenbeck & Ronquist, 2001), but with different branch lengths.

REFERENCES

- Althoff, D.M., Gitzendanner, M.A. & Segraves, K.A. (2007) The utility of amplified fragment length polymorphisms in phylogenetics: A comparison of homology within and between genomes. *Systematic Biology* **56**, 477-484.
- Braun, V., Buchner, O. & Neuner, G. (2002) Thermotolerance of photosystem 2 of three alpine plant species under field conditions. *Photosynthesis* **40**, 587-595.
- Doyle, J. & Doyle, J. (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* **19**, 11-15.
- Huelsenbeck, J.P. & Ronquist, F. (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**, 754-755.
- Hunt, R. (1982) *Plant growth curves: the functional approach to plant growth analysis*. Edward Arnold, London.
- Swofford, D.L. (1999) PAUP: phylogenetic analysis using parsimony (*and other methods). Sinauer Associates, Inc., Sunderland, MA.
- Vos, P., Hogers, R., Bleeker, M., Reijnders, M., Van De Lee, T., Hornes, M., Frijers, A., Pot, J., Peleman, J., Kuiper, M. & Zabeau, M. (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* **23**, 4407-4414.
- Xia, C. & Zhenghai, H. (2000) Comparative studies on leaf structure and oil cells of the Magnaliaeae in China. *Acta Phytotaxonomica Sinica* **38**, 218-230.

```
#NEXUS
BEGIN D101;
dimensions ntax=10 nchar=219;
format missing=?
symbols="ABCDEFGHIJKLMNPQRSTUVWXYZ"
interleave datatype=DN1 gap= -;
```

```
matrix
grandisA      1100000111001100??00100000000000100011000001100001
hookeriA      1101001111000011??11000000000001110011000001101111
insignis2     11011101111111110??00001110000011110011100001111111
szechuanica   1101000111000110??01000110000110110011100001101101
Liriodendron  1101110111000110??0000000000000110011000001100001
megaphyllaA   0101110111111111??10101100000001110011100001100001
kwangtungensisA 0101111111000110??0010110000000110011000001100001
fordianaA     1111000111000111??11001010001100100011000001100001
chingji2      010100011100001000100001101??001100011000001101111
pachyphyllaA  1111110111000111??10000110010001100011000001111111
```

```
grandisA      1111111111001??11011110000100110100110001100101000
hookeriA      11110000000010011011100001100??0100110011100100001
insignis2     1111011111??11011001010000111111100110000111110001
szechuanica   0111010111??10111001010000001111000110010111111101
Liriodendron  1011011111??10011011110000001111100110000100100001
megaphyllaA   1011011111??11011011110000111111100111101111111111
kwangtungensisA 1111011111??11011011011000111111100110001100110001
fordianaA     1111011111??101111101100001111110001111111111110001
chingji2      1111111110??00011010110000001000000110001000100000
pachyphyllaA  1111111111??01111010111000111001001111111100100110
```

```
grandisA      1111001??10010111111111011111100110110110001111100
hookeriA      1100001?11101011111111?011111110100111110001101101
insignis2     1111001??1000001101111101111111111011110001111111
szechuanica   1111001????????????????????????????????????????????
Liriodendron  1111001??00000011111111010111000001110110000111111
megaphyllaA   1111111??10011011111111111011100111111110010111111
kwangtungensisA 1111001??11100011111111011110101100111110011111111
fordianaA     1111111??0000001111111001111010011111110001101101
chingji2      0111001??10001011001111010001110100011110001101101
pachyphyllaA  1111001??10001011111111000001000100010011101101101
```

```
grandisA      100001111?110011111111111111011100110110111111110
hookeriA      101111111?1100111111111101111101100110110110111110
insignis2     101011111?110011111000111011010000110111111111110
szechuanica   ??????????11111111101111101101000011101111011110
Liriodendron  101111111?110010110000100001010000110000110011110
megaphyllaA   101111111?111111110110101001111010110110111111110
kwangtungensisA 101111111?1111001110001110110110001101101110011110
fordianaA     110110011?111011110111111110000001101001111111111
chingji2      100110011????????????????????????????????????????????
pachyphyllaA  100110011?1100001100111111000100000100101100111111
```

```
grandisA      1011001111110011111
hookeriA      1111111111110011101
insignis2     1111111011110110111
szechuanica   1110111000011010111
Liriodendron  1110111000010000111
megaphyllaA   1110111011110001111
kwangtungensisA 1110111011110011111
fordianaA     1101111010111010111
chingii2      ???????????????????
pachyphyllaA  1111111010110110111
;
end;
```

species	Amax	LSP	LCP	Thickness of leaves	Thickness of palisade tissue
Mg	6.72335	723.928	13.8522	324	73
Mh	5.7779	693.28	11.4931	291	65
Mf	9.00883	896.851	8.25178	282	155
Mc	8.1938	893.155	10.5395	288	70
Mp	9.07798	935.542	10.9794	297	79
Mi	9.2088	751.654	7.86899	340	93
Ms	10.2165	896.145	10.3506	313	96
Mm	8.76503	713.161	10.1885	206	43
Mk	8.08355	755.73	11.1994	293	84

photosynthetic

leaf structure

Thickness of sponge tissue	Basal stem diameter	Crown volume	Leaf area	Relative growth rate
186	8.6525	51083.796	852.025	5.9530922
161	7.91	39736.33	993.27	7.4094938
58	7.58	41989.981	838.7725	6.1240256
166	8.93	133913.27	1582.96	8.4508945
167	9.04	48037.927	836.71	7.7592505
162	9.77	49741.544	1006.8267	7.0802655
198	8.4425	26817.085	803.42	5.6839331
98	8.5575	33772.291	1152.0695	7.2950712
150	8.95	61019.91	1217.195	7.2849033

plant growth

Tch	Tmax	T50
45.98	52.471	45.7225
43	50.63175	44.985
43.795	50.72925	45.965
45.025	50.66325	46.875
44.585	52.01975	46.76
41.34	49.3665	43.675
43.2275	51.66275	45.53
44.75	51.283	46.2675
46.025	51.91425	47.675

Photosynthetic thermal tolerance

APPENDIX S4: DERIVATION FOR THE STATISTICAL METHODS USED TO ESTIMATE SIMULTANEOUSLY TRAIT CORRELATIONS AND PHYLOGENETIC SIGNAL

The model underlying our statistical methods assumes that the evolution of two or more traits follows an Ornstein-Uhlenbeck (OU) process of stabilizing selection. For simplicity, we give the derivation in terms of two traits (x and y) measured for N species, with the extension to multiple traits obvious. Also, we describe the complete model including phylogenetic signal and measurement error; the special cases with no measurement error, or analyses of single traits separately, can be derived by excluding the appropriate components of the model.

MODEL OF THE EVOLUTIONARY PROCESS

We assume that both traits x and y have a single optimum for all species, yet the strength of selection towards these two optima can differ between species; this formulation is a 2-trait extension of the single-trait OU process as modelled by Blomberg *et al.* (2003). Furthermore, we assume that evolution of the two traits is correlated with correlation coefficient r . Over some arbitrarily small time step Δt , the evolution of trait values $X(t)$ and $Y(t)$ for a given species is described by

$$X(t+\Delta t) = d_x X(t) + \gamma_x(t)$$

$$Y(t+\Delta t) = d_y Y(t) + \gamma_y(t) \tag{eqn S1}$$

where d_x and d_y measure the strength of stabilizing selection for traits x and y , and $\gamma_x(t)$ and $\gamma_y(t)$ are random variables with means zero, variances σ_x^2 and σ_y^2 , and correlation r . Under the

assumption that the variance in the value of a trait for the ancestral species at the base of the tree is zero, the covariance in trait x between species i and j is

$$\text{cov}\{X_i, X_j\} = d_x^{\tau_i + \tau_j} \frac{1 - d_x^{2\tau_{ij}}}{1 - d_x^2} \sigma_x^2 \quad \text{eqn S2}$$

where X_i and X_j are the values of trait x for species i and j , τ_i and τ_j are the distances from the node representing their most recent common ancestor to tip of the tree for species i and j , and τ_{ij} is the branch length from the base to their most recent common ancestor (Blomberg *et al.*, 2003). A similar expression holds for the covariance in values of trait y among species. Using a similar approach, it can be shown that the covariance between trait x for species i and trait y for species j is given by

$$\text{cov}\{X_i, Y_j\} = d_x^{\tau_i} d_y^{\tau_j} \frac{1 - (d_x d_y)^{\tau_{ij}}}{1 - d_x d_y} r \sigma_x \sigma_y \quad \text{eqn S3}$$

Note that when $d_x = d_y = 1$, the covariance matrix derived directly from the model of Brownian motion evolution is recovered. When $d_x = 0$, the covariances in trait x among species become zero, and the variances in trait x for each species is σ_x^2 ; this holds similarly for trait y .

Our model is similar to previous models that lead to phylogenetic signal. Felsenstein (1988) was the first to suggest using an OU process for describing stabilizing selection, while Garland *et al.* (1993) developed the first statistical implementation of the OU process for simulation-based hypothesis testing using the PDSIMUL program. Martins and Hansen (1997) and Hansen (1997) were the first to develop likelihood-based statistical approaches based on the

OU process. In contrast to the formulation of Martins and Hansen (1997), who assume that the trait values for the basal species on the phylogenetic tree are taken from the stationary distribution of the OU process, we follow Blomberg et al. (2003) in assuming that the variance in the trait values at the base of the tree is zero. This has the practical advantage that, as d approaches one, the covariances among species approach those expected under a Brownian motion (BM) model of evolution. Because the BM model is a random walk model in continuous time, it is a non-stationary process and therefore has no stationary distribution. In addition to capturing a BM process as a special case, our formulation also allows for phylogenetic signal stronger than given by a BM process ($d > 1$). Although Martins and Hansen (1997) and Hansen (1997) use the OU model to estimate phylogenetic signal, they do not consider other types of analyses, such as correlation or regression. Butler & King (2004; , 2005) and Scales *et al.* (2009) extend the methods of Hansen (1997) and provide computer code to allow the finer-scale analysis of evolutionary patterns of single traits within phylogenetic trees.

Our approach is also similar to that of Pagel (1994), who derives a test for the correlated evolution of two binary traits that each take only two values. His test is based on a model of evolution in which trait values switch between states (0 and 1) as species evolve along a phylogenetic tree, with the rate of switching determining the phylogenetic signal. This is analogous to the model we describe above, in which continuous-valued traits evolve along a phylogenetic tree. There are numerous, similar approaches for binary traits (Grafen & Ridley, 1997; Schluter, Price, Mooers *et al.*, 1997; Lorch & Eadie, 1999; Mooers & Schluter, 1999; Pagel & Meade, 2006). A related approach for analyzing binary traits is given by Ives and Garland (Ives & Garland, accepted), who develop a method of phylogenetic logistic regression that incorporates an estimated parameter for phylogenetic signal.

Using Pagel's λ (1997), Freckleton et al. (2002) develop methods for both univariate analyses for the detection of phylogenetic signal in a single trait, and correlation analyses that include Pagel's λ as a simultaneously estimated measure of phylogenetic signal. This approach is not based upon an evolutionary model; Pagel's λ is simply a discounting parameter that is multiplied to phylogenetic covariances among species. Unlike a model based on a specific evolutionary process, there is no guarantee that the resulting statistical model is well-formulated. Specifically, it is possible for estimation of λ to lead to infeasible covariance matrices that are not positive definite; for our model developed around an evolutionary model, this cannot occur. Furthermore, λ is generally constrained to be less than one, making the analyses incapable of revealing phylogenetic patterns that are stronger than those produced by BM evolution.

Our methods for performing correlation analyses while estimating phylogenetic signal are similar to those we have developed for regression analyses. Lavin et al. (2008) present methods and Matlab code to perform regression under the assumption that the dependent variable evolves according to an OU process, and also implements methods using Pagel's λ (Pagel, 1997), Grafen's rho (Grafen, 1989), and the ACDC model of Blomberg *et al.* (2003).

Finally, there are several other, more distantly related methods for measuring phylogenetic signal (Cheverud, Dow & Leutenegger, 1985; Diniz-Filho, de Sant'Ana & Bini, 1998; Abouheif, 1999). These methods are generally inferior to model-based methods that allow more rigorous statistical tests and evolutionary inferences, but a thorough comparison is beyond the scope of this paper (see also Rohlf, 2006).

STATISTICAL MODEL

The general structure of the statistical model is

$$\mathbf{X}^* = a_x + \boldsymbol{\varepsilon}_x(d_x)$$

$$\mathbf{Y}^* = a_y + \boldsymbol{\varepsilon}_y(d_y) \tag{eqn S4}$$

$$\mathbf{X} = \mathbf{X}^* + \boldsymbol{\eta}_x$$

$$\mathbf{Y} = \mathbf{Y}^* + \boldsymbol{\eta}_y$$

where \mathbf{X}^* and \mathbf{Y}^* represent the $N \times 1$ vectors containing true values of traits x and y among species, a_x and a_y are the means of traits x and y , $\boldsymbol{\varepsilon}_x(d_x)$ and $\boldsymbol{\varepsilon}_y(d_y)$ give the variability in traits x and y that may contain phylogenetic signal measured by the parameters d_x and d_y , and \mathbf{X} and \mathbf{Y} are the values observed with measurement error (within-species variation) incorporated with the random variables $\boldsymbol{\eta}_x$ and $\boldsymbol{\eta}_y$. Assuming that evolution follows an OU process, the covariance matrix for $\boldsymbol{\varepsilon}_x(d_x)$, namely $\text{cov}\{\mathbf{X}^*, \mathbf{X}^*\} = \sigma_x^2 \mathbf{C}_x(d_x)$, has elements given by equation S2, and $\mathbf{C}_y(d_y)$ is defined similarly for $\boldsymbol{\varepsilon}_y(d_y)$. The covariance matrix across traits x and y , $\text{cov}\{\mathbf{X}^*, \mathbf{Y}^*\} = r\sigma_x\sigma_y\mathbf{C}_{xy}(d_x, d_y)$, has elements given by equation S3.

The observed values of traits x and y can be expressed in terms of both $\boldsymbol{\varepsilon}$ and $\boldsymbol{\eta}$ as

$$\mathbf{W} = \mathbf{A} + \boldsymbol{\varepsilon} + \boldsymbol{\eta} \tag{eqn S5}$$

where \mathbf{W} is the $2N \times 1$ vector created by stacking \mathbf{X} and \mathbf{Y} , and \mathbf{A} is the $2N \times 1$ vector whose first N elements are a_x and second N elements are a_y , and $\boldsymbol{\varepsilon}$ and $\boldsymbol{\eta}$ are $2N \times 1$ vectors created by stacking $\boldsymbol{\varepsilon}_x(d_x)$ and $\boldsymbol{\varepsilon}_y(d_y)$, and $\boldsymbol{\eta}_x$ and $\boldsymbol{\eta}_y$, respectively. The resulting covariance matrix

$$\text{cov}\{\mathbf{W}, \mathbf{W}\} = \sigma^2 \boldsymbol{\Psi} \text{ is}$$

$$\sigma^2\Psi = \begin{pmatrix} \sigma_x^2\mathbf{C}_x(d_x) + \sigma_{mx}^2\mathbf{M}_x & r\sigma_x\sigma_y\mathbf{C}_{xy}(d_x, d_y) + \sigma_{mx}\sigma_{my}\mathbf{M}_{xy} \\ r\sigma_x\sigma_y\mathbf{C}_{xy}(d_x, d_y) + \sigma_{mx}\sigma_{my}\mathbf{M}_{xy}' & \sigma_y^2\mathbf{C}_y(d_y) + \sigma_{my}^2\mathbf{M}_y \end{pmatrix}, \quad \text{eqn S6}$$

where $\sigma_{mx}^2\mathbf{M}_x$ and $\sigma_{my}^2\mathbf{M}_y$ are matrices containing the within-species trait covariances, and $\sigma_{mx}\sigma_{my}\mathbf{M}_{xy}$ contains the covariances in measurements among traits.

The key feature of the joint covariance matrix (eqn S6) is that it contains covariances among species in the matrices \mathbf{C} that reflect phylogenetic relationships (and phylogenetic signal d) and correlation between traits given by r . Although the correlation coefficient r is often thought of as a measure computed directly from the data, here it is better considered as a parameter in a statistical model given by the covariance matrix. The estimates of r , and the estimates of d_x and d_y , are those values that provide the best fit of the covariance matrix to the data.

STATISTICAL MODEL FITTING

Following Ives et al. (2007), we assume that magnitudes of measurement error variances are known. Estimates of the measurement error variances ($\sigma_{mx}^2\mathbf{M}_x$ and $\sigma_{my}^2\mathbf{M}_y$) can be obtained from the variation in the trait values among individuals within each species. For example, the matrix $\sigma_{mx}^2\mathbf{M}_x$ contains the square of the standard errors for values of trait x for each of the N species as diagonal elements, and all non-diagonal elements are zero. When data for different traits are collected using different individuals from the same species, then there should be no covariance between the measurements of different trait values. However, in our data set different traits were measured on the same plants, and therefore the measurement error of values

for different traits may be correlated. We can incorporate this covariance into $\sigma_{mx}\sigma_{my}\mathbf{M}_{xy}$ that contains as diagonal elements the covariances in measurement errors of the values of traits x and y measured for each of the N species; the off-diagonal elements are zero. If there are n individuals of species i , then the covariance in measurement errors (the i -th diagonal element of $\sigma_{mx}\sigma_{my}\mathbf{M}_{xy}$) is $\frac{1}{n(n-1)}\sum_{k=1}^n(x_{i(k)} - \bar{x}_i)(y_{i(k)} - \bar{y}_i)$ where $x_{i(k)}$ and $y_{i(k)}$ are the values of x and y for the k -th individual of species i .

The parameters in the model that must be estimated are d_x and d_y for phylogenetic signal, r for the correlation between trait values, and σ_x^2 and σ_y^2 for the among-species variance in trait values. We used REML estimation which involves minimizing the conditional log-likelihood function in the estimated parameters d , r , and σ^2 (for a technical discussion, see Ives *et al.*, 2007). For small data sets, the restricted likelihood function may not be unimodal. Therefore, we used simulated annealing as a robust approach to minimization, followed by Nelder-Mead minimization to polish the parameter estimates. These analyses are performed by the Matlab code CORRELATIONv2.m that is available from TG upon request.

REFERENCES

- Abouheif, E. (1999) A method for testing the assumption of phylogenetic independence in comparative data. *Evolutionary Ecology Research* **1**, 895-909.
- Blomberg, S.P., Garland, T., Jr. & Ives, A.R. (2003) Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* **57**, 717-745.
- Butler, M.A. & King, A.A. (2004) Phylogenetic comparative analysis: A modeling approach for adaptive evolution. *American Naturalist* **164**, 683-695.

- Cheverud, J.M., Dow, M.M. & Leutenegger, W. (1985) The quantitative assessment of phylogenetic constraints in comparative analyses - sexual dimorphism in body-weight among primates. *Evolution* **39**, 1335-1351.
- Diniz-Filho, J.A., De Sant'ana, C.E.R. & Bini, L.M. (1998) An eigenvector method for estimating phylogenetic inertia. *Evolution* **52**, 1247-1262.
- Evans, M.E.K., Hearn, D.J., Hahn, W.J., Spangle, J.M. & Venable, D.L. (2005) Climate and life-history evolution in evening primroses (Oenothera, Onagraceae): A phylogenetic comparative analysis. *Evolution* **59**, 1914-1927.
- Felsenstein, J. (1988) Phylogenies and quantitative characters. *Annual Review of Ecology and Systematics* **19**, 445-471.
- Freckleton, R.P., Harvey, P.H. & Pagel, M. (2002) Phylogenetic analysis and comparative data: a test and review of evidence. *American Naturalist* **160**, 712-726.
- Garland, T., Jr., Dickerman, A.W., Janis, C.M. & Jones, J.A. (1993) Phylogenetic analysis of covariance by computer-simulation. *Systematic Biology* **42**, 265-292.
- Grafen, A. (1989) The phylogenetic regression. *Transactions of the Royal Society of London B, Biological Sciences* **326**, 119-157.
- Grafen, A. & Ridley, M. (1997) A new model for discrete character evolution. *Journal of Theoretical Biology* **184**, 7-14.
- Hansen, T.F. (1997) Stabilizing selection and the comparative analysis of adaptation. *Evolution* **51**, 1341-1351.
- Ives, A.R. & Garland, T., Jr. (accepted) Phylogenetic logistic regression for binary dependent variables. *Systematic Biology*.

- Ives, A.R., Midford, P.E. & Garland, T., Jr. (2007) Within-species variation and measurement error in phylogenetic comparative methods. *Systematic Biology* **56**, 252-270.
- Lavin, S.R., Karasov, W.H., Ives, A.R., Middleton, K.M. & Garland, T., Jr. (2008) Morphometrics of the avian small intestine, compared with non-flying mammals: a phylogenetic approach. *Physiological and Biochemical Zoology* **81**, 526-550.
- Lorch, P.D. & Eadie, J.M. (1999) Power of the concentrated changes test for correlated evolution. *Systematic Biology* **48**, 170-191.
- Martins, E.P. & Hansen, T.F. (1997) Phylogenies and the comparative method: a general approach to incorporating phylogenetic information into the analysis of interspecific data. *American Naturalist* **149**, 646-667. Erratum 153:448.
- Mooers, A.O. & Schluter, D. (1999) Reconstructing ancestor states with maximum likelihood: Support for one- and two-rate models. *Systematic Biology* **48**, 623-633.
- Pagel, M. (1994) Detecting correlated evolution on phylogenies: a general method for the comparative analysis of discrete characters. *Proceedings of the Royal Society of London Series B-Biological Sciences* **255**, 37-45.
- Pagel, M. (1997) Inferring evolutionary processes from phylogenies. *Zoologica Scripta* **26**, 331-348.
- Pagel, M. & Meade, A. (2006) Bayesian analysis of correlated evolution of discrete characters by reversible-jump Markov chain Monte Carlo. *American Naturalist* **167**, 808-825.
- Rohlf, F.J. (2006) A comment on phylogenetic correction. *Evolution*. **60**, 1509-1515.
- Scales, J.A., King, A.A. & Butler, M.A. (2009) Running for your life or running for your dinner: what drives fiber-type evolution in lizard locomotor muscles? *American Naturalist* **173**, 543-553.

Schluter, D., Price, T., Mooers, A.O. & Ludwig, D. (1997) Likelihood of ancestor states in adaptive radiation. *Evolution* **51**, 1699-1711.